

The Environment of Al Asfar Evaporation Lake in Al Hassa, Saudi Arabia: An Integrated Study

A.S. El Mahmoudi, Y.Y. Al-Dakheel, A.H.A. Hussein and M.A. Massoud
Water Studies Center, King Faisal University, Al Hofuf, KSA

Abstract

There is a growing concern in the Kingdom of Saudi Arabia to exploit water resources of acceptable quality to meet demands in domestic, industrial, and agricultural sectors. Treated wastewater and desalinated water are among the various alternative supplies to meet these demands.

Al Hassa Oasis is one of the main and old agricultural centers in Saudi Arabia. An irrigation network of concrete canals and drainage network were put into operation in 1971. The project delivers 328 million m³ of spring water for about 22, 000 farms. The excess drainage water is collected by earthen drainage network and discharged into two evaporation lakes which are Al Asfar and Al Uyoun.

Such Evaporation lakes (Wet lands) are important link between water and land ecosystems. They provide habitats and refuges for wildlife, flood protection and contribute to abatement of impact of pollution and promoting groundwater recharge. The main salient morphologic features of Al Asfar Lake are wet lands, sabkhas and sand dunes. There are salt tolerant vegetation (halophyte) found in some of the less salt affected sabkha areas.

The present study aims to study the environment of Evaporation Al Asfar Lake, Al Hassa area. Remote sensing, sedimentological, hydrogeological, chemical studies and Geographical Information Systems (GIS) are implemented to achieve the objectives of this study.

The processed landsat-5 TM Imageries indicated, there is a general increment in both the sabkha and water ponds areas extent with the passage of time. The origin and the various sediments types and their statistical grain size parameters were elaborated. Through chemical analyses, the major and minor ions content of soil pore water, surface water and sabkha soil samples were assessed. The results of this study are important in any future development of this area of the Al Asfar and in planning of how to reuse the water lakes in the agricultural purposes.

Keywords: Evaporation Lakes, Inland Sabkha, Water Reuse, Arid Environment, Remote Sensing, GIS, Soil Chemistry, Water Quality, Al Hassa Oasis, Saudi Arabia.

Introduction

Water resources are essential and play a significant role in the development processes. The KSA is known by its arid conditions and limited renewable freshwater resources. Surface water in KSA is very limited and of a little significance in the water budget of the KSA. To meet the increasing water demands in domestic, industrial, and agricultural sectors, various alternative supplies are surface water, renewable groundwater resources, treated wastewater, desalinated water, and non-renewable groundwater.

Fresh water supply in arid regions is limited due to scarce precipitation and high evapotranspiration. When the limited supply is coupled with increasing demand for fresh water in these regions due to rapid population growth, it necessitates optimal development and management of water resources.

Al Hassa Oasis is one of the main and old agricultural centers in Saudi Arabia. Al Hassa oasis with an area of 20,000 hectares is located 70 km west of the Arabian Gulf (Figure. 1). An irrigation network of 1, 450 Km of concrete canals and drainage network were put into operation in 1971 (Abderrhman & Bader, 1992). The project delivers 328 million m³ of spring water for about 22, 000 farms. Nowadays this amount of water is increased due to use of the treated wastewater from Al Hofouf sewage station.

Al Hassa drainage water is collected by 1641 earthen lateral open canals having depths of about 1 m. The collected drainage water in the eastern sector of the project flows from the laterals to three sub main drainage canals which are connected to main drainage canal D2 (Figure 2). The collected water in this drainage flows towards an evaporation lake at 13 Km (Figure 2) to the east of Al Hassa Oasis. The drainage water in the northern sector flows from the laterals to six sub main drainage canals which are connected to main drainage canal D1. Drainage water in D1 flows towards an evaporation lake outside Al Hassa Oasis, 52 km north of Al Hofuf. The water remains in the evaporation lakes during winter season to be used to supply cattle with drinking water and for aquifer recharge. From December until April of each year, the agricultural drainage water overflows from the northern lake through small waterways into the Arabian Gulf, (Abderrhman & Bader, 1992).

The main salient morphologic features of Al Asfar Lake are wet lands, sabkhas and sand dunes. There are salt tolerant vegetation (halophyte) found in some of the less salt affected sabkha areas. The area is characterized by widespread growth of halophyte shrubs associated with a very thin salt crust on the sabkha surface (Figure 3). In desert environments sabkha is one of the least investigated environments. That is most probably due to its subtle nature. However, due to its potential economic value and/or its direct or indirect impact on

the environment; particularly groundwater, interest in the sabkha deposits increased in recent years. Information on the location and area of various grades of sabkha will provide valuable information about these phenomena and to know its environmental impacts on the surrounding areas and to investigate its dynamics.

The Eastern province of Saudi Arabia has many sabkha; the coastal sabkha has been studied by many workers (e.g., Barth and Boer, 2002, .Mah, 2004 and Barth, 2006).

A very distinct vegetation type which occurs at both inland and coastal sabkhat is the *Phoenixdactylifera* / *Tamarix* sp type. These sites are highly valued by recreation seeking people because they provide shade and green in the otherwise unspectacular landscape. Generally there are much more locations along the sabkha edges where the environmental conditions would support Phoenix/Tamarix communities. It is strongly suggested that these communities be cultivated in order to provide more recreational possibilities especially nearurban settlements (Barth, 2006).

Hydrogeology of Al Hassa area have been reported in a number of studies (BRGM, 1977 and 1981, Etewy et al. 1983, HIDA, 1986, Al-Mahmoud, 1987, Abdurrahman, 1988, Edgell, 1989, Al Dakheel & Al Safarjalani, 2005 and Al Safarjalani and Almadini 2007 etc.).

The present study aims to study the environment of Evaporation Lake of Al Asfar, Al Hassa area. Remote sensing, sedimentological, hydrogeological, chemical studies and Geographical Information Systems (GIS) are implemented to achieve the objectives of this study. The main concern of this paper is the presentation of some quantitative data on the geochemistry of sabkha soil water and on the geochemistry of the solid constituents of the crustal layer of Al Asfar sabkha and ascertain the maturity class of Al Asfar sabkha in reference to the classification proposed by Bahafzallah et al. (1993).

Materials and Methods

Multi-date landsat-5 TM imageries of Years 1987, 1993, 1998 and 2000 of the study area have been processed and analyzed to investigate the dynamics of both water ponds and the associated sabkha with time at Al Asfar lake area. One hundred and eight surface soil samples representing each soil series from Al Asfar sabkha were collected. The sampling of sabkha was sometimes hindered by small water bonds or local vegetations. Fourteen samples from surface water and the isolated water bonds and twenty samples of soil water (from the area characterized by shallow water table (≈ 20 cm depth) have been collected in May to June, 2007. The latitude and longitude of the soil and water samples coordinates was measured by using high accuracy differential GPS (Trimble ProXR GPS). The collected soil samples were raised from pits 20-30 cm deep.

The soil samples were then air dried, grounded and sieved through a 2 mm sieve. A paste in deionized water was prepared and filtered. Extracts of soil samples are prepared and tagged for sedimentological & chemical analysis. The grain size has been done using the Granular Composition Test Set with sieve shaker and standard set of sieves for dry and wet sieving manufactured by Eijkelkamp. The granular composition of the representative sample obtained by applying an electromagnetic sieve shaker. The sieve shaker keeps the sample continuously in motion in order to obtain the best possible sieving results. The pH and total soluble salts were measured in the soil paste extract (Rhoades, 1982). Sodium and potassium was determined by flame photometry according to Jackson (1973). Calcium and magnesium were determined by atomic absorption spectrophotometer according to Carter (1993). Soluble carbonates and bicarbonates were determined volumetrically in the soil paste extract by titration against 0.01 N Hydrochloric acid using phenolphthalein and methyl orange as indicators according to Jackson (1967). Soluble chlorides were determined by titration with 0.01 N silver nitrate solution and potassium chromate as indicator according to Richards (1954). Organic matter content was determined according to Walkley-Black rapid titration method (Jackson, 1967). The concentrations of NO_3^- were determined in soil paste extract and water samples according to Norman et al. (1985) method. ArcGIS software and satellite image processing software (ERDAS Imagine) were used for the integration of the output parameters of this study and for visualizes their spatial distribution.

Results and Discussions

A) Remote sensing Investigations

In this study, Landsat -5 TM imageries with seven bands of Years 1987, 1993, 1998 and 2000 of the study area have been geo-referenced, visually interpreted and analyzed to identify different geomorphic units of Al Asfar Lake environment. The digital elevation model (DEM) of the study area was also processed and analyzed. The Elevation Data collected using the Trimble GPS system has been used for constructing a Digital Elevation Model (DEM) for Al Asfar area (Figure 4). This map facilitated the interpretation of the result of the grain size distribution and the result of the chemical analysis. The processed landsat-5 TM Imageries indicated, there is a general increment in both the sabkha and wet land areas extent with the passage of time (sabkha & wetlands in 1987) vs. (sabkha & wetlands in 2000), (Figure 5). The direction of increment is related to relief where increased sabkha areas have taken place in low relief areas where groundwater rose and pushed salt to the surface. Sabkha spectral signature in imagery taken at different wavelengths is complex. It is related to salt content in soil, low relief, moisture content, groundwater table,

surface water in sabkha areas, salt tolerant halophytic vegetation, texture, color and composition of soil and soil profiles. However, different sabkha grades are recognized. These different sabkha grades have various degrees of salt content as it is indicated from the TDS spatial distribution map gained from the chemical analysis task of this study.

B) Sedimentological Results

Forty one representative samples out of the 108 collected sabkha soil samples, were selected and the following analyses were carried out: grain size analysis, determination of carbonate, sand and mud contents and organic matter content. In the following is the discussion of the result of each analysis.

Grain Size Analysis

The main purpose of grain size analysis is to get an idea about the size spectrum of the sabkha samples under investigation. It also gives information on the energy of the medium of deposition, where coarse particles are usually transported in higher energy environment, whereas, and low energy environment transports finer sediments. It is also worthy to add that the statistical parameters of grain size and their mutual relationships provide us with information on agents of transportation as well as the environment of deposition.

Grain size statistical parameters form the basis of many schemes for classifying sedimentary environments. The size distribution of coarse clastic sediments reflects the fluidity (viscosity) factor of the depositing medium and the energy factor of the environment (site) of deposition (Sahu 1964). Grain size parameters include graphic mean size (Mz), inclusive graphic standard deviation (σ_I), inclusive graphic skewness (SK_I) and graphic kurtosis (K_G) are calculated. The estimation of the statistical grain-size parameters was accomplished using the mathematic formulas of Folk & Ward (1957). Table (1) shows the descriptive statistics of the different grain size parameters for the analyzed sabkha soil samples. Surfer contouring software has been used to draw spatial distribution maps for these parameters, these maps will generally used to estimate the relation between grain size distribution and the sedimentological and geomorphologic features of the area.

Table 1: Descriptive statistics of the different grain size parameters for the analyzed sabkha soil samples, Al Asfar lake area, Al Hassa, KSA.

Variable	Mz (ϕ)	σ_I (ϕ)	SK _I	K _G
Range	0.81:2.31	0.56:1.53	-0.22:- 0.28	0.73:- 1.27
Average	1.50	0.96	0.08	1.03
Standard Deviation	0.31	0.24	0.12	0.12

The average value of Mz is 1.5Φ (medium sand). The Mz values indicate that most of the samples of sabkha deposits are composed of medium sand. Figure (6, a) represents the spatial distribution of Mz for the analyzed samples in the study area. This map indicates that the fining is getting towards the north and to the west of the area of study. This distribution is related to the sources of sediments and the geomorphologic features including the drainage system and the topography of the study area (see Fig. 4). The dominance of the fine sediments at the western part of the study area are coming from the mouth of the main Drain (D2) and the floods of the main lake of Al Asfar.

The average value of sorting (σ) is 0.96Φ (moderately sorted). The samples of poorly sorted are located to the west of the study area Figure (6, b), this may be attributed to the different sources of the sediments having different sediment characteristics (i.e. sabkha, alluvial deposits and aeolian sediments, and dust storms) and the effect of geomorphology. While moderately sorted and moderately well sorted are located to the east of the study area.

The average values of SK_i is 0.08Φ (near symmetrical). The data obtained showed that the most sediment of all the studied sabkha deposits are near symmetrical in the middle and to the north of the area while the southern part samples are fine skewed Figure (6, c). It is worth to mention that there is a good match between the map of Mz distribution (Fig.6 a) and the map of SK_i (Fig. 6 c), see for example the patch off the negative skewed around the location of samples 6 and 7 which means fine particles and that is clearly indicated in the map of Mz (Fig. 6, a). Also, the southern part of the map is characterized by positive values of SK_i which means coarse particles good match with map of Mz (Fig. 6, a). Areas with positively skewed sediments are formed in association with the tidal flats and sabkhas, (Al Hurban et al., 2007).

The sediment samples in the study area are inclined towards the "mesokurtic" (average value of kurtosis is kG 1.03Φ , ranging between platykurtic (0.73) to leptokurtic (1.29), Figure (6, d). The data obtained for KG indicates that the Al Asfar sabkha sediments are mesokurtic (60.97 %), leptokurtic (24.39 %) and platykurtic (14.63 %). This may also be attributed to the variety of sediment sources and the complexity of surface geomorphology of the area. Platykurtic sediments are mostly associated with the poorly sorted sediments and this is clear if you compare the western part (poorly sorting) of map shown in figure (6,b) with the western part (platykurtic)of map shown in Figure (6,d).

In general, the grain size statistical analyses indicate that most of the areas are composed of two or more classes of sediments transported and deposited from different sources; these are aeolian sand, sabkhas and drain input and coastal features. The variability in the grain size statistical parameters may be attributed to the complexity of surface morphology as well as the diversity in the type of depositional environment in Al Asfar area.

Determination of Carbonate, Sand and Mud contents

Figures 7, a, b & c show the spatial distribution of the carbonate, sand and mud respectively. From the grain size analysis and from these maps, it is clear the Al Asfar sabkha are in general sand and sandy silt. Also, It could be noticed the clearly that both carbonate and mud contents are generally increase in the west of the area of study and are attributed to the discharged water from the main drain network, mainly D-2 (see Fig. 2).

Organic matter content

The organic matter content for these sabkha soil samples is analyzed. The high values of the organic matter content are belonging to samples located quite near to the periphery of the main lake of Al Asfar (Figures 7, d) and this is attributed to the accumulation of the decayed vegetation with depth at these locations.

C) Chemical and Hydrochemical Investigations

The samples were analyzed for their physical properties; hydrogen ion concentration (pH), electrical conductivity (E.C.), and then total dissolved salts (T.D.S.) is calculated and for major constituents (Na^+ , Ca^{++} , K^+ , Mg^{++} , Cl^- , SO_4^{--} , HCO_3^- and NO_3^-). Table (2) summarizes the statistical analysis of physical properties and for major constituents of the collected sabkha soil, surface water and pore water samples. From this table it could be noticed, the major cations are in order Na^+ , K^+ , Ca^{++} and Mg^{++} , while the major anions are in order: Cl^- , SO_4^{--} , HCO_3^- and NO_3^- . The carbonate content is so small that it was undetected as the maximum value of pH is not more than 8.3 which is the threshold this ions start to form for both soil and water samples (Stumm and Morgan, 1981).

Table (2): Statistical analysis for the parameters of physical properties (pH, Ec, TDS) and for the major cations and anions for Al Asfar sabkha soil (108 samples), surface water and for pore water Samples (34 samples).

Type	Analysis Values	pH	Ec	TDS	Major Cations				Major Anions			
					Na^+	Ca^{++}	K^+	Mg^{++}	Cl^-	So^-	Hco_3^-	No_3^-
Soil Samples	Minimum	7.22	5.6	4464.0	561.0	188.2	229.2	92.7	1014.6	718.7	31.1	2.5
	Maximum	8.28	70.9	56720.0	7262.1	2308.3	2775.4	1178.2	15146.1	12891.7	863.3	95.0
	Mean	7.653	27.7	22166.1	2841.3	900.7	1085.9	459.0	6130.7	4243.9	112.2	12.1
	SD*	0.188	15.8	12635.7	1619.1	513.9	618.7	261.8	3513.6	2578.3	82.3	13.0
Water Samples	Minimum	7.0	9.5	7624.0	1008.9	235.8	159.7	536.3	1827.2	1349.9	70.0	0.9
	Maximum	8.3	102.3	81840.0	10829.6	2531.5	1714.3	5756.7	27335.0	8809.8	1742.4	41.0
	Mean	7.8	40.5	32383.5	4285.2	1001.7	678.4	2277.9	10717.7	4321.9	318.7	9.1
	SD*	0.3	28.7	22973.4	3040.0	710.6	481.2	1616.0	8079.9	2447.0	287.0	9.4

*Standard Deviation

PH was measured for all sediment samples and water samples. Figure 8, a & b shows the spatial distribution map of pH values for soils and water along the study area. The soil samples having pH values varying between 7.22 and 8.28,

while the water samples having pH values 7 and 8.3. From these two maps, it could notice that pH Values are increasing southwestward in opposite trend of TDS distribution map (Fig. 9 a & b). Obviously all the samples indicate a slightly alkaline character with very narrow pH ranges.

For the studied sabkha sediments, the total salinity ranges between 4,464 and 56,720 ppm, with mean value of 22166.1 ppm. The increasing of soil and water salinity towards the east (Figure 9, a) is quite expected due to the effect of leaching process in the down-gradient direction. The lower gradient in salinity content towards the southwestern part can be attributed to the flushing effect of drain two (D-2) while the higher salinity content recorded for the zone located at the eastern side of the main lake of Al Asfar (Figure 9, a) and have the same trend of the lake extension (NW-SE). The higher salinity content is attributed to the influence of the intermittent charge and for the high evaporation rate. From the spatial distribution of salinity content of the sabkha soils, it is clear that salt content has a significant effect on the sabkha grade and has its effect on sabkha spectral signature in landsat-5 TM imageries used in this study (Figure 5).

The total salinity of surface water and pore water samples ranged from 7624 to 81840 ppm with mean value of 32383.5 ppm (Figure 9, b). The relatively low values of total salinity are belonging to the surface water samples from the main lake of Al Asfar (Figure 9, b). This relatively low salinity values encourage to re-treat Al Asfar lake water and to reuse it in agricultural purposes, this will lead to minimize the sabkha extension, to mitigate its impact on the nearby farms of Al Hassa oasis and to mitigate aquifer recharge with high saline water and to optimize the water use of this surface water body in this arid environment. The increase of salinity of this lake is due to its closed nature, shallow depth of water, restricted condition and minor connection with continuous fresh water source.

The soil water salinity at the margins of Al Asfar Lake ranges from less than 10,000 ppm (see for example salinity of samples Nos. Als1, Als5 & Als7) to about 80,000 ppm (see for example salinity of samples Nos. Als 6, Als11 & Als19), (Figure 9, b). The salt content of soil water for the collected samples depends on the distance from the main water body of Al Asfar lake, i.e. the nearest to the lake, the less of salt content. The soil water under this sabkha is more saline than the normal sea water.

It is worthy to mention that the contour pattern of the constructed spatial distribution maps for the major cations of soil samples (Figure 10, a) are similar to that of TDS shown in Figure (9 a). The same feature is also noticed for the major cations of water samples (see Figure 9, b & Figure 11, b). However, in case of anions the contour pattern for chloride and sulphate have the similar pattern of TDS while for both Bicarbonate and Nitrate have a different pattern Figures (11 a & 11 b). The bicarbonate in this sabkha is thought to be derived partly from the outcrops of Neogene carbonate ridges originated under marine conditions. It is worthy to mention that the bicarbonate ions distribution either in soil or in water (Figure 11, a & b) is in a good match with carbonate content distribution of soil sabkha samples shown in (Figure 7, a).

The major hydrodynamic process operating in this inland sabkha is upward migration of the brines by capillary movement precipitating salts. The presence of salts at the surface due to the rate of evaporation at the sabkha surface at these sites exceeds than the total input of water to the sabkha, so that high concentration of TDS in the water are reached (Handford, 1981). The variation in the salinity is relatively influenced by the following factors; evaporation, the lithologic characteristics of the soils, as well as transportation, solubility of salts and the rate of discharge of the surface water from the agricultural drains to Al Asfar Lake.

The rate of evaporation in the inland sabkhas is supposedly higher than that in the coastal ones due to the more arid conditions. Consequently, the ground-water table plays a substantial role in the development of inland sabkhas, which are usually less developed than coastal sabkha flats and are predominantly tectonically and/or topographically controlled (El-Naggar, 1988). The sediments of these sabkhas consist predominantly of gypsum (desert roses), quartz and calcite, with halite always existing at the crust (Kinsman, 1969). The crustal part is the product of a combination of geological processes like desiccation, weathering and, above all, cementation of the salts precipitated through evaporation of highly saline groundwater. Khedr (1989) terms such a growth or development of sabkhas through this latter process as "sabkhaization".

Further, these results shed light on the maturity status of the inland Sabkha of Al Asfar. Thus the observed higher TDS in both soil and water confirms the slightly developed to developed nature of sabkha as per the classification of Bahafzallah et al. (1993).

Conclusions

Integration of the processed Landsat TM imagery, sedimentological results & chemical analysis contributes to a better understanding of the environment of Al Asfar Lake, Al Hassa area. The processed landsat-5 TM Imageries indicated there is a general increment in both sabkha and wet land areas extent with the passage of time.

Chemical analyses of the surface water of Evaporation lake of Al Asfar and soil water and the sabkha soil are interpreted. Different sabkha grades are recognized. These different sabkha grades have various degrees of salt content as it is indicated from the TDS spatial distribution at Al Asfar area. This study revealed that the surface water of Al Asfar lake is characterized by a relatively low salinity and then re-treating and reusing of this water in agricultural purposes will lead to minimize the sabkha extension, to mitigate its impact on the nearby farms of Al Hassa oasis, to mitigate aquifer recharge with high saline water and to optimize the water use of this available surface water body in this arid environment.

Chemical results are very important to be taken in consideration by Saudi Aramco, as the different sabkha grades have their various corrosion levels that may impact the pipelines crossing this sabkha area. Sabkha dynamics information and chemical results gained from this study may be used to optimize pipelines

maintenance and future cost effective pipelines route planning. The evaporation condition prevailing in the present study and the absence of continuous recharge from the main drainage (D2) led to the concentration of highly saline lagoon water. The major hydrodynamic process is possibly the upward migration subsurface of brines from groundwater by capillary action due to evaporation, precipitating salt on the surface.

Acknowledgments

The authors would like to express their sincere appreciation to Deanship of Scientific Research, King Faisal University for the financial support of this study under fund grant No. 8095. Special thanks go to Mohamed Al Kulaib and Mohamed Al Farj for their assistance in the field activities and in the laboratory measurements of this study.

References

- Abderrahman, W. A and Bader, T.A., (1992). Remote Sensing use in the management of Agricultural Drainage Water in Al Hassa Oasis of Saudi Arabia, Remote Sensing in Evaluation and Management of Irrigation. Remote Sensing Env. 237-246.
- Abdurrahman, W., (1988). Water management plan for the Al-Hassa Irrigation and Drainage Project in Saudi Arabia. Agric. Water Manage., 13, 185-194.
- Al Dakheel, Y.Y and A.M., Al Safarjalani, (2005). Study of the historical pattern of changes in water quality of Al Hassa Oasis springs. J. of Productivity and Development, 10 (2), 211-225 (in Arabic).
- Al Hurban, A., El-Gamily, H. and El-Sammak, A., (2007). Geomorphic changes in Ras Al-Subiyah area, Kuwait. Environmental Geology, DOI, 10.1007/s00254-007-0919-z.
- Al-Mahmoud, M.J. (1987). Hydrogeology of Al Hassa Oasis, M.Sc. Thesis, King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia, 34 p.
- Al Safarjalani, A. M. and A.M. Almadini (2007). Effect of Spatial Variations in Chemical Properties of Irrigation Groundwater at Al-Hassa Oasis, Kingdom of Saudi Arabia, Damascus University ,Damascus University Journal for Agricultural and Scientific Research , 23 (1),141-168.

- Bahafzallah A., Laila A., Kazi A. and Al-saify M., (1993). Classification and distribution of the Red Sea coastal sabkhas near Jeddah- Saudi Arabia. *J. Carbonates and Evaporites*, Vol. 8, (1), 23-28.
- Barth, H.-J. & Böer, B. (2002) (Eds.), *Sabkha Ecosystems, The Arabian Peninsula and Adjacent Countries*. Dordrecht, Netherlands: Kluwer.
- Barth, H.-J. (2006). Sabkha edge vegetation of coastal and inland sabkhat in Saudi Arabia. In: Khan, M.A. & Weber, D.J. (Hrsg.) *Ecophysiology of High Salinity Tolerant Plants*, Springer., 215-225.
- BRGM, Bureau de Recherches Geologiques et Ministeres, (1977). *Al Hassa Development Project Groundwaters Study and Management Programme. Final Report vol. 1, Groundwater Resources Development Department, Ministry of Agriculture and Water, Riyadh, KSA.*
- BRGM, Bureau de Recherches Geologiques et Minieres, (1981). *Reutilization of Drainage Water in Al Hassa Oasis*". Ministry of Agriculture and Water, Riyadh, Saudi Arabia, 2, pp. 3 and 161.
- Carter, M.R. (1993). *Soil Sampling and Methods of Analysis*. Canada Society of Soil Science, Lewis Publishers.
- Edgell, H.S., (1989). Geological framework of Saudi Arabia groundwater resources, *Journal of King Abdulaziz University, Earth Sciences*, Vol.3. Scientific Publishing Centre, Jeddah, 267-286.
- El-Naggar, Z.R. (1988). *Foundation Problems in Sabkha Deposits. Short Course: Foundation Engineering for Practicing Engineer, KFUPM, Dhahran, April, SD1-SD54.*
- Etewy, H., M. Assed, S. Al-Barrak and A. M. Turjoman, (1983): Water quality and soil characteristics as related to irrigation and drainage in Al-Hassa area. A paper presented in "The Sixth Conference on the Biological Aspects of Saudi Arabia" Held in College of Science, King Abdulaziz University, Jeddah, KSA, March 1-3, 1983. 489-811.
- Folk, R.L. and Ward, W.C. (1957). Brazos River bar: a study in the significance of grain size parameters. *Journal of Sedimentary Petrology*, 27(1), 3-26.
- Handford, C. R., (1981). A process-sedimentary framework for characterizing recent and ancient sabkhas: *Sedimentary Geology*, v. 30, 255–265.
- HIDA, Hassa Irrigation and Drainage Authority, (1986). *The Effect of salinity in Groundwater and Surface water on the Cultivated Crops in Al Hassa Oasis*".

Symposium on the Effect of Water Quality on Human Health and Agriculture,
Organized by SWICC and GCC, Al-Khobar, Saudi Arabia.

Jackson, M.L. (1967). Soil Chemical Analysis. Prentice Hall, Inc., Engle Wood Cliff.,
USA.

Jackson, M.L. (1973). Soil Chemical Analysis. Prentice-Hall, Inc. India.

Khedr E.S., (1989). Recent coastal sabkhas from the Red Sea: a model of
sabkhaization. Egypt. J. Geol., Vol. 33, No. 1-2, 87-120.

Kinsman, D.J.J. (1969). Modes of Formation, Sedimentary Associations, and
Diagnostic Features of Shallow-Water and Supratidal Evaporates. Reprinted
from: Amer. Assoc. Petroleum Geologists Bull., v. 53, pp. 830-840, published
in: Marine Evaporites: Origin, Diagenesis, and Geochemistry. Edited
by: Douglas W. Kirkland and Robert Evans, Dowden, Hutchinson & Ross, Inc.,
pp. 80-90.

Mah, A. (2004). A remote sensing study of sabkha in the eastern province of Saudi
Arabia, Proceedings of the Third International Conference & Exhibition, GIS-
2004, 27-29 September, 2004, Kingdom of Bahrain, 263-280.

Norman, R.J., J. C. Edberg and J. W. Stucki. (1985). Determination of nitrate and
nitrite in soil extracts by ultraviolet spectrophotometry. Soil Sci. Soc. Am., J.
49:1182-1185.

Rhoades, J.D. (1982). Soluble salts P. 167-185. In A.L. Page et al. (ed.) Methods
of soil analysis Part 2. Chemical and microbial methods. . 2nd ed. Agron.
Monogr 9. ASA and SSA. Madison, WI.

Richards, L.A. (ed) (1954). Diagnosis and improvement of saline and alkali soils.
U.S. Salinity laboratory staff. Agr. Hand book No. 60.

Sahu, BK (1964). Depositional mechanisms from the size analysis of clastic
sediments. J Sed Petrol 34(1), 73-83.

Stumm W. and Morgan J.J. (1981). Aquatic Chemistry: An Introduction
Emphasizing Chemical Equilibria in Natural Waters. J. Willey, New York, 583
p.




Figure 1: Base map of Al Hasa Region and the location of the study area.




Figure 2: Main Drainage System of Al Hasa Irrigation and Drainage Project.




Figure 3: Different types of salt tolerant vegetation (halophyte) at Al Asfar area.






Figure 4: Digital Elevation Model Map of Al Asfar area, Al Hassa, KSA.




a) Landsat-5 TM, October, 1987




b) Landsat-5 TM, November 2000


Figure 5: The processed landsat-5 TM Imageries for Al Asfar area.




a) Mz



b) σ



c) SKI



d) KG

Figure 6: Distribution of Grain Size Statistical Parameters in the Study Area, a) Mean Grain Size, b) Sorting, c) Skewness and d) Kurtosis.




Figure 7: The spatial distribution of carbonate, sand, mud and organic matter contents in the inland sabkha samples of Al Asfar area, Al Hassa, KSA.




Figure 8: Distribution map of the pH content for the soil and water of Al Asfar inland Sabkha, Al Hassa area, KSA.




Figure 9: Spatial Distribution map of the total salinity a) for the soil, b) for the surface water and soil water of Al Asfar area, Al Hassa area, KSA.




Figure 10, a: Distribution maps of the major cations concentration for the soil of Al Asfar area, Al Hassa, KSA.




Figure 10, b: Distribution maps of the major cations concentration for the surface water and soil water of Al Asfar area, Al Hassa, KSA.




Figure 11, a: Distribution maps of the major anions concentration for the soil of Al Asfar area, Al Hassa, KSA.




Figure 11, b: Distribution maps of the major anions concentration for the surface water and soil water of Al Asfar area, Al Hassa, KSA.